

READ THE LABEL BEFORE USE

Gladiator

160 me

Reg. No./Nr. L9846 Act/Wet 36/1947

A systemic micro-emulsion herbicide for the control of woody plants and weeds as listed for forestry, grass pastures, conservation and industrial areas.

'n Sistemiese emulgeerbare konsentraat onkruidodder vir die beheer van houtagtige plante en onkruid soos aangetoon in bosbou, grasweidings en nywerheidsgebiede.



DANGER

Hazard statements

- Causes serious eye irritation
- Causes serious eye damage
- Very toxic to aquatic life

Precautionary statements

- Wear protective gloves/ protective clothing/ eye protection/ face protection

HERBICIDE GROUP CODE	O	ONKRUIDDODER GROEPPKODE
----------------------	---	-------------------------

ACTIVE INGREDIENT

Picloram (Pyridine Compound)
(as tri-isopropanolamine salt)
Fluroxypyr (Pyridine Compound)
(as methyl heptyl ester)

/
80 g/l
80 g/l

AKTIEWE BESTANDDEEL

Pikloram (Piridienverbinding)
(as tri-isopropanolamiensout)
Fluroksipir (Piridienverbinding)
(as metiel-heptielester)

NET VOLUME (5 ℓ)

REGISTRATION HOLDER/REGISTRASIE HOUER:

Future Farm & Forest
Services & Supplies (Pty) Ltd.
Reg. 1998/014639/07
P.O. Box 98165
Sloane Park
2152



ECOGUARD

Environmentally compatible solutions for
pest & vegetation management

NETTO VOLUME (5 ℓ)

DISTRIBUTED BY/ VERSPREI DEUR:

Ecoguard Biosciences (Pty) Ltd.
Reg. 1988/001616/07
Tel: 011 - 463 6057
Fax: 011 - 463 6462

UN NUMBER
BATCH NUMBER
DATE OF MANUFACTURE

3082

UNNUMMER
LOTNUMMER
DATUM VAN VERVAARDIGING

EMERGENCY CONTACT Tel: 011 463 5842
TYGERBERG POISON INFO CENTRE: 086 155 5777

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

WARNINGS:

- Poisonous by swallowing.
- Causes serious eye damage or serious eye irritation.
- May causes allergic skin reactions.
- Store in a cool place.
- Very toxic to aquatic life.
- Toxic to aquatic life with long lasting effects.
- Store away from food, feed, seed and other agricultural chemicals and out of reach of children, uninformed persons and animals.

RE-ENTRY INTERVAL:

Do not enter treated area until spray deposit has dried on the target plants unless wearing protective clothing.

GRAZING RESTRICTIONS:

Meat animals grazing up to two weeks after treatment should be removed from treated area three days prior to slaughter. Do not graze lactating dairy animals on treated area within two weeks after treatment.

GRAZING POISONOUS PLANTS:

Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of weeds against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration.

Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Obtain, read, and follow all safety instructions before use.
- Wear protective gloves and face shield or safety glasses when handling the concentrate.
- Do not breathe fumes or spray mist.

- Wash contaminated clothing daily.
- Wash hands thoroughly after handling with soap and water.
- Avoid eye contact. In case of eye contact, flush eyes with clean water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- If eye irritation persists: Get medical help.
- If exposed or concerned: Call a POISON CENTER/doctor.
- Do not eat, drink or smoke whilst applying, mixing or before washing hands and face and change of clothing.
- Prevent spray drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the spray applicator with a 1 % solution of household ammonia before using with other products. Allow the solution to stand in the spray tank for several hours but preferably overnight. Rinse at least twice with clean water. This applicator should not be used for applying products other than herbicides. Dispose of wash water where it will not contaminate food, grazing, rivers or dams.
- Rinse empty containers three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsing to the contents of the spray tank before destroying the container by perforation and flattening. NEVER use for other purposes.
- Prevent contamination of food, feeds, drinking water and eating utensils.
- Avoid release to the environment. – if this is not the intended use.
- Collect spillage.
- Store locked up.
- Dispose of contents/container in accordance with applicable regulations.

RELEVANT SUBSTANCES

	Weight (%)
Picloram triisopropanolamine salt	12-16%
Fluroxypyr 1-methylheptyl ester	8-12%

SYMPTOMS OF HUMAN POISONING:

There are no specific signs of poisoning. The concentrate may cause severe eye irritation. Prolonged or repeated skin contact may cause allergic reactions in some individuals. Harmful effects not anticipated from swallowing small amount. Prolonged exposure to vapours not likely to be hazardous.

FIRST AID TREATMENT:

EYES: Irrigate with clean flowing water for 30 minutes. Seek medical advice.

SKIN: Remove contaminated clothing. Rinse affected area thoroughly with clean running water.

INGESTION: In case of accidental ingestion, do NOT induce vomiting. Give 1 or 2 glasses of water or milk to drink. Do not give anything by mouth to an unconscious person. Seek medical attention.

INHALATION: Remove patient to fresh air. If effects occur, seek medical advises.

NOTE TO PHYSICIAN:

No specific antidote. Treatment should always be symptomatic. If lavage is performed, suggest endotracheal and / or esophagoscopy control.

USE RESTRICTIONS:

Precautions for avoiding injury to non-target plants are:

- Do not permit spray mist to drift or come into contact with sensitive broadleaf crops, including, but not limited to lucerne, beans, melons, potatoes, soyabeans, sunflowers, tobacco, tomatoes, cotton, fruit trees, grape vines, ornamentals, soils containing roots of these plants, soil in which such plants are to be grown, grain varieties in a susceptible stage of growth or grazing or any other area not under treatment.
- Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray or spray drift or spray run off to fall onto banks or bottoms of irrigation ditches, canals, streams, dams, rivers, either dry or containing water that may be used for irrigation or domestic purposes or may carry water to an irrigation facility.
- Do not make application when circumstance favour movement or drift from the treated site.
- Do not apply to areas that may be rotate to any broadleaf crop.
- Do not use manure from animals grazing treated areas on land for growing broadleaf crops, ornamentals, orchards or other susceptible crops. Manure may contain enough picloram to cause injury to susceptible plants.
- Do not use grass or sprayed plants from treated areas for composting or mulching of susceptible broadleaf plants or crops.
- Do not transfer livestock from treated grazing areas (or feeding of treated grass) onto sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated grass pasture. Otherwise, urine and manure may contain enough picloram to cause injury to broadleaf plants.
- Do not use on flood irrigated land or fields.
- Do not spray if the loss of forage legumes, including clover cannot be tolerated. New legume seedlings may take years to re-establish.
- Do not move treated soil to areas other than sites for which **GLADIATOR 160 ME** is registered for use.
- Do not apply to land near to desirable broadleaf plants or land onto which such plants are to be grown or grain varieties in a susceptible stage of growth. The product may however be applied in the vicinity of desirable broadleaf plants, or soil or land onto which such plants are to be grown, or grain varieties in a susceptible stage of growth, provided that adequate precautionary measures are taken to avoid spray drift or contamination of run-off areas.
- Apply the product strictly in accordance with the application directions.

DIRECTIONS FOR USE: Use only as directed.

COMPATIBILITY:

GLADIATOR 160 ME is compatible with ACTIPRON SUPER**.

MIXING:

For foliar applications, half fill the spray tank with water and add the required quantity of **GLADIATOR 160 ME**. While filling the spray tank add ACTIPRON SUPER at the rate indicated below while maintaining constant agitation.

APPLICATION:

A. FOLIAR APPLICATION

- Use only on actively growing plants with full leaf cover.
- Apply as a full cover spray to leaves and stems to the point of spray solution run-off.
- Do not apply to foliage wet from rain or dew etc.
- Avoid spray drift onto non-target plants.
- Avoid fine droplet size. Use a knapsack sprayer with a solid cone nozzle (eg. Spraying systems TG-1, or equivalent type) that will ensure an even coverage of the target area.
- Maintain a constant pressure of between 200 – 250 kPa. Maintain approximately 50 cm from the target for best coverage.
- Rain 3 hours after application will have no effect on efficacy provided the spray has dried on the target plants.
- Prevent drift of spray mist onto desirable plants.

B. CUT STUMP

- Apply preferably immediately after felling but definitely within 1 hour.
- **GLADIATOR 160 ME** should not be applied to the stump if surface is wet from rain or dew.
- Apply as a coarse spray using solid cone nozzle tips (eg. Spraying Systems TG-1 or equivalent type) at 100 kPa, low pressure.
- Do not exceed spray height of 20 cm above target. Do not apply if wind velocity exceeds 15 km/h (as measured by handheld wind recorder). The difference between the wet and dry bulb readings on a whirling hygrometer must not exceed 8 °C.
- Apply according to directions in the table below.

Warning:

If top kill is obtained but regrowth occurs from the crown, apply an herbicide registered for use against the relevant species as a foliar spray.

RATES AND METHODS:

A. FOLIAR APPLICATIONS:

NOTE: ACTIPRON SUPER should be added at a rate of 500 ml / 100 litres spray mix for all foliar applications.

PLANT	RATE / 100ℓ WATER	REMARKS
<i>Acacia mearnsii</i> (Black Wattle)	125 ml (0.125% mixture)	See DIRECTIONS FOR USE above. Apply to actively growing plants. Apply until the point of run-off.
<i>Acacia longifolia</i> (Long leaf wattle)	500 ml (0.50% mixture)	
<i>Acacia melanoxylon</i> (Blackwood)		
<i>Acacia saligna</i> (Port Jackson)		
<i>Anredera baselloides</i> (Madeira vine)		
<i>Campuloclinium macrocephalum</i> (Pom pom weed)	375 ml (0.375% mixture)	Plants too high for a good cover spray should be slashed and the re-growth sprayed when not less than 0,5 m high.
<i>Chromolaena odorata</i> (Triffid weed)	750 ml (0.75% mixture)	
<i>Cirsium vulgare</i> (Scotch thistle)		
<i>Eucalyptus grandis</i> (Saligna gum)	500 ml (0.50% mixture)	In the event of any re-growth after application, a follow-up spray should be carried out.
<i>Lantana camara</i> (Lantana)	1,5 ℓ (1.50% mixture)	
<i>Psidium guajava</i> (Guava)		
<i>Robinia pseudoacacia</i> (Black Locust)		
<i>Rubus cuneifolius</i> (American bramble)	750 ml (0.75% mixture)	
<i>Solanum mauritianum</i> (Bugweed)	250 ml (0.25% mixture)	

B. CUT STUMP

NOTE: ACTIPRON SUPER should be added to the spray mixture at a rate of 500 ml / 100 litres for all cut stump applications.

PLANT	RATE / 100ℓ WATER	REMARKS
<i>Acacia mearnsii</i> (Black Wattle)	2.0 ℓ (2.00% mixture)	See DIRECTIONS FOR USE above. Apply to low cut stumps (10 – 20 cm high) preferably with a single cut surface. Apply to complete cut surface and root crown of stumps with a diameter of less than 10 cm. Where multiple stumps are present, all cut surfaces must be treated. For bigger stumps, apply to the cambial region (sapwood) of the cut surface and root crown.
<i>Acacia melanoxylon</i> (Blackwood)		
<i>Acacia saligna</i> (Port Jackson)		
<i>Chromolaena odorata</i> (Triffid weed)	750 ml (0.75% mixture)	
<i>Eucalyptus cladocalyx</i> (Sugar gum)	4.5 ℓ (4.50% mixture)	
<i>Eucalyptus grandis</i> (Saligna gum)	3.0 ℓ (3.00% mixture)	
<i>Eucalyptus smithii</i> (Blackbutt peppermint)	4.5 ℓ (4.50% mixture)	
<i>Lantana camara</i> (Lantana)	1.5 ℓ (1.50% mixture)	
<i>Melia azedarach</i> (Syringa)		
<i>Ricinus communis</i> (Castor Oil Plant)	1.0 ℓ (1.00% mixture)	In all cases, apply until the point of runoff. In the event of some re-growth and depending on the species, a follow-up spray as a coppice application.

In situations where more than one species of target plant occurs contact your local technical representative.

For assistance in the calibration for spot treatments please contact your local technical representative.

ACTIPRON SUPER** (Mineral oil), Reg. No. L5506, Act No. 36 of 1947, is the Trademark of H&R GSP (Pty) Ltd.

WAARSKUWINGS:

- Giftig indien ingesluk word.
- Konsentraat mag ernstige irritasie van oë veroorsaak.
- Mag allergiese velreaksie veroorsaak.
- Effens giftig vir waterlewendende invertebrata.
- Berg in 'n koel plek.
- Baie giftig vir waterlewe.
- Giftig vir waterlewe met langdurige effekte.
- Berg weg van voedselware, voer, saad en ander landboumiddels en buite die bereik van kinders, oningeligte persone en diere.

HERBETREDINGSINTERVAL:

Moenie behandelde area binnegaan voordat die spuitstof op behandelde plante afgedroog het nie, tensy beskermende klere gedra word nie.

WEIDINGSBEPERKINGS:

Vir die eerste twee weke na toediening moet slagdiere minstens 3 dae voor slag uit behandelde area onttrek word. Moenie diere in laktasie vir melkproduksie binne twee weke na behandeling in behandelde area laat wei nie.

WEI VAN TOKSIESE PLANTE:

Onkruidodder behandeling mag die smaaklikheid van sekere toksiese plante verhoog. Moenie behandelde areas laat beweï voordat toksiese plante dood en dus nie meer smaaklik vir diere is nie.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande; kwaliteit van verdunningswater; verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die onkruid teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings van die betrokke middel na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie.

Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMATREËLS:

- Verkry, lees en volg alle veiligheidsinstruksies voor gebruik.
- Dra beskermende handskoene en gesigskerm of veiligheidsbril wanneer die konsentraat hanteer word.
- Moenie dampe of sproeiereën inasem nie.
- Was besoedelde klere daaglik.
- Was onmiddellik met seep en water indien per ongeluk met die vel in aanraking kom.
- Vermoeg aanraking met die oë. In geval van oogkontak, spoel oë met skoon water.
- INDIEN IN OË: Spoel versigtig uit met water vir 'n paar minute. Verwyder kontakklense, indien teenwoordig en maklik om te doen. Gaan voort met spoel.
- Indien oogirritasie voortduur: Kry mediese hulp.
- Indien blootgestel of bekommerd: Ontbied 'n GIFSENTRUM/dokter.
- Moenie eet, drink of rook tydens vermenging en toediening en alvorens hande en gesig gewas en skoon klere aangetrek is nie.
- Verhoed spuitstof wegdrywing na ander gewasse, weidings, riviere, damme of enige gebied wat nie behandel word nie.
- Maak spuittoerusting met 'n 1 % huishoudelike ammoniakoplossing skoon voordat ander plaagdoders daarmee gebruik word. Laat staan die oplossing in die spuittenk vir 'n paar uur, maar verkieslik oornag. Spoel ten minste twee keer met skoon water uit. Hierdie spuittoerusting behoort nie gebruik te word vir die toediening van enige ander chemikalieë nie, behalwe onkruidodders. Gooi waswater weg waar dit nie voedsel, weiding, riviere of damme sal besoedel nie.
- Spoel leë houer driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10 % van die houer. Gooi die spoelwater by die inhoud van die spuittenk voordat die houer vernietig word deur gate daarin te maak en dit plat te druk en moet nie vir enige ander doel gebruik nie.
- Voorkom besoedeling van voedselware, voer, drinkwater en eetgerei.
- Vermoeg vrystelling na die omgewing. – indien dit nie die beoogde gebruik is nie.
- Versamel gemors.
- Stoor toegesluit.
- Gooi inhoud/houer weg in ooreenstemming met toepaslike regulasies.

RELEVANTE STOWWE

Pikloram tri-isopropanolamien sout
Fluroksipir 1-metiel-heptielester

Inhoud (%)

12-16%
8-12%

SIMPTOME VAN VERGIFTIGING:

Daar is geen spesifieke tekens van vergiftiging nie. Die konsentraat mag ernstige oog irritasie veroorsaak. Verlengde en aanhoudende velkontak mag allergiese reaksies in sommige individue veroorsaak. Skadelike effekte van die opneming van klein hoeveelhede is nie verwag nie. Lang blootstelling aan dampe sal nie noodwendig gevaarlik wees nie.

NOODHULPBEHANDELING:

OË: Spoel uit met skoon lopende water vir 30 minute. Verkry mediese advies.

VEL: Verwyder die besoedelde klere. Spoel besmette liggaamsdele deeglike met skoon lopende water.

INDIEN INGESLUK: In geval van toevallige inname, gee een of twee glase water of melk om te drink. Moet nie die persoon laat braak nie. Nul per mond vir 'n bewustelose persoon. Verkry mediese advies.

INDIEN INGEASEM: Verwyder die persoon na 'n plek met vars lug. Verkry mediese advise.

NOTA AAN GENEESHEER:

Geen spesifieke teenmiddel. Behandeling simptome. Indien maagspoeling gedoen moet word, word endotracheale en/of esofagoskopie aanbeveel.

GEBRUIKSBEPERKINGS:

Maatreëls om beskadiging van nie-teiken plante te voorkom is soos volg:

- Voorkom dat spuitnewel wegdryf of in kontak kom met sensitiewe breëblaar gewasse, insluitend, maar nie beperk tot: lusern, bone, waatlemoene, aartappels, sojabone, sonneblomme, tabak, tamaties, katoen, vrugtebome, wingerdstokke, sierplante, gronde
- waarin wortels van die plante voorkom, grond waarin die plante verbou gaan word, jong graangewasse in vatbare groeistadium, weidings of enige ander area wat nie behandel gaan word nie.
- Moenie water vir besproeiing of huishoudelike gebruik besoedel nie. Om beskadiging van gewasse of ander gewenste plante te voorkom, moenie toelaat dat spuitnewel of spuitstof op walle of bodem van besproeiingsvore, kanale, strome, damme, riviere, wat
- vir besproeiing of huishoudelike gebruik bedoel is, hetsy droog of met water daarin, beland of deur water na besproeiings sisteme afgevoer word nie.
- Moenie onder toestande wat die beweging van die onkruid doder uit behandelde areas bevorder, toedien nie.
- Moenie areas behandel wat moontlik met enige breëblaar gewas geroteer gaan word nie.
- Moenie die mis van diere wat op behandelde areas gewei het vir lande waarop 'n breëblaar gewas, sierplante, boorde of enige ander vatbare gewas gevestig is of gevestig gaan word, gebruik nie. Mis mag genoeg onkruid doder nog bevat om skade te veroorsaak op gevoelige gewasse.
- Moenie gras of bespuite plante uit behandelde areas vir kompostering of deklaag by gevoelige breëblaar plante of gewasse gebruik nie.
- Moenie lewende hawe van behandelde weidingsareas (of wat op behandelde gras gevoed het) na areas van gevoelige breëblaar gewasse oorplaas alvorens hulle vir minstens 7 dae op onbehandelde grasweidings gewei het nie. Urine of mis van sulke diere mag voldoende picloram bevat om gevoelige breëblaar plante te beskadig.

- Moenie op lande of veld wat vloedsbesproei word gebruik nie.
- Moenie toedien as die verlies van peulgewas weidings, insluitend klawers, nie geduld kan word nie. Dit mag jare neem voordat nuwe peulplante sal vestig. Moenie behandelde grond na areas anders as gronde waarop die gebruik van **GLADIATOR 160 ME** geregistreer is, verskuif nie. Moenie op grond wat NABY gewenste breëblaar plante of grond waarop sulke plante groei, of graangewasse wat in gevoelige ontwikkelings stadium is toedien, mits daar nie voldoende voorsorg getref word om die wegdrywing van spuitnewel na die areas of besoedeling van afloop areas te voorkom nie.
- Dien die produk streng volgens die gebruiksaanwysings toe.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

MENGBAARHEID:

GLADIATOR 160 ME is mengbaar met ACTIPRON SUPER**.

VERMENGING:

Vir looftoediening vul die spuittenk halfvol met water en voeg die benodigde hoeveelheid **GLADIATOR 160 ME** by. Vul die spuittenk dan verder met water en voeg ACTIPRON SUPER by teen die dosis soos aangedui hieronder, terwyl roering plaasvind.

TOEDIENING:

A. LOOFTOEDIENING:

Wend slegs aan wanneer plante volle drag blare het en aktief groei.

- Dien toe as 'n volledige dekbespuiting op loof en stamme tot die punt van spuitstof afloop.
- Moet nie toegedien word wanneer blare nat is as gevolg van reën of dou ens.nie.
- Vermyn spuitstofwegdrywing op nie-teiken plante.
- Vermyn fyn druppelgroottes.
- Gebruik 'n rugsakspuit wat toegerus is met soliede keelspuitstukke (bv. Spraying Systems TG-1, of soortgelyke spuitstukke) wat egalige benatting sal verseker.
- Handhaaf 'n konstante druk van 200 – 250 kPa. Handhaaf 'n afstand van ongeveer 50 cm van die teiken vir gewenste benatting.
- Na toediening sal reën 3 ure na toediening geen invloed op die effektiwiteit hê nie, indien die spuitstof op die teiken plant gedroog het.
- Verhoed spuitingsstofwegdrywing na gewenste plante.

B. GEKAPTE STOMPESPUITING:

- Dien toe verkieslik onmiddellik na afkap maar definitief binne 1 uur.
- **GLADIATOR 160 ME** moet nie toegedien word op stompe indien hulle nat is weens reën of dou nie.

- Dien toe as 'n growwe spuit deur gebruik van soliede keelspuitstukke (bv. Spraying Systems TG-1, of soortgelyke spuitstukke) en handhaaf 'n druk van 100 kPa, (lae druk).
- 'n Spuihoogte van 20 cm bokant die teiken moet nie oorskry word nie. Indien die windspoed hoër is as 15 km/u (soos bepaal deur 'n handwindmeter) moet die produk nie toegedien word nie. Die verskil tussen die nat- en droëbollesing, soos met 'n swaaihigrometer bepaal moet nie meer as 8 °C wees nie.
- Dien toe volgens aanwysings vir stompe en gekapte stompbespuiting in tabel hieronder.

A. LOOFTOEDIENNING

OPMERKING: ACTIPRON SUPER, moet ten alle tye vir alle loofbespuitings bygevoeg word teen 500 ml / 100-liter finale spuitmengsel.

PLANTSOORT	DOSIS / 100ℓ WATER	OPMERKINGS
<i>Acacia mearnsii</i> (Swartwattel)	125 ml (0.125% mengsel)	Sien GEBRUIKSAANWYSINGS hierbo. Dien toe op jong aktief groeïende plante tot die punt van afloop.
<i>Acacia longifolia</i> (Langblaarwattel)	500 ml (0.50% mengsel)	
<i>Acacia melanoxylon</i> (Swarthout)		
<i>Acacia saligna</i> (Port Jackson)		
<i>Anredera baselloides</i> (Madeira vine)		
<i>Campuloclinium macrocephalum</i> (Pom pom onkruid)	375 ml (0.375% mengsel)	Plante te hoog vir goeie spuitbedekking moet eers gekap word. Die hergroeï word gespuit wanneer dit nie minder as 0,5 m hoog is nie.
<i>Chromolaena odorata</i> (Paraffienbos)	750 ml (0.75% mengsel)	
<i>Cirsium vulgare</i> (Skotsedissel)		
<i>Eucalyptus grandis</i> (Salignabloekom)	500 ml (0.50% mengsel)	In die geval van enige hergroeï na bespuiting, moet 'n opvolg bespuiting toegedien word.
<i>Lantana camara</i> (Lantana)	1,5 ℓ (1.50% mengsel)	
<i>Psidium guajava</i> (Guava)		
<i>Robinia pseudoacacia</i> (Witakasia)		

<i>Rubus cuneifolius</i> (Amerikaanse braam)	750 ml (0.75% mengsel)	
<i>Solanum mauritianum</i> (Luisboom)	250 ml (0.25% mengsel)	

B. GEKAPTE STOMPESPUITING

OPMERKING: ACTIPRON SUPER, moet ten alle tye vir alle stompbespuitings bygevoeg word teen 500 ml / 100-liter finale spuitmengsel.

PLANTSOORT	DOSIS / 100ℓ WATER	OPMERKINGS
<i>Acacia mearnsii</i> (Swartwattel)	2.0 ℓ (2.00% mengsel)	Sien GEBRUIKSAANWYSINGS hierbo.
<i>Acacia melanoxylon</i> (Swarthout)		
<i>Acacia saligna</i> (Port Jackson)		
<i>Chromolaena odorata</i> (Parriënbos)	750 mℓ (0.75% mengsel)	Dien toe op laag gekapte stompe (10 – 20 cm hoog) verkieslik met 'n enkele gekapte oppervlakte. Dien toe op die hele gekapte oppervlakte en wortelkroon vir stompe met 'n deursnee van minder as 10 cm. Waar veelvoudige stompe teenwoordig is, moet alle gekapte oppervlakte behandel word.
<i>Eucalyptus cladocalyx</i> (Suikerbloekom)	4.5 ℓ (4.50% mengsel)	
<i>Eucalyptus grandis</i> (Salignabloekom)	3.0 ℓ (3.00% mengsel)	
<i>Eucalyptus smithii</i> (Swartstampeperment)	4.5 ℓ (4.50% mengsel)	
<i>Lantana camara</i> (Lantana)	1.5 ℓ (1.50% mengsel)	
<i>Melia azedarach</i> (Sering)		
<i>Ricinus communis</i> (Kastorolie plant)	1.0 ℓ (1.00% mengsel)	Vir groter stompe, dien toe op die cambium gedeelte (splinthout) en wortelkroon. In alle gevalle, dien toe tot die punt van afloop. In die geval van hergroei mag 'n opvolgbespuiting, afhangend van die spesie, benodig word.

GLADIATOR 160 ME beheer bome en struie. Ander bome en struie wat nie tydens die ontwikkelingsproewe met die middel voorgekom het nie, kan ook in meerdere of mindere mate daardeur beheer word. Die registrasiehouer aanvaar egter geen aanspreeklikheid ten opsigte van ongelyste bome en struie nie.

BELANGRIK: Wanneer meer as een ongewenste plant spesies voorkom, raadpleeg die plaaslike tegniese verteenwoordiger van die verskaffer.

Vir hulp met kalibrasie by kolbehandeling raadpleeg die plaaslike tegniese verteenwoordiger van die verskaffer.

ACTIPRON SUPER** (mineraal olie) (Reg. Nr. L5506, Wet Nr. 36 van 1947), is die handelsmerk van H&R GSP (Edms) Bpk.